



# PennState Health

## Emergency Medicine Residency Program Director

Penn State Health Milton S. Hershey Medical Center is seeking an Emergency Medicine Residency Program Director to join our exceptional academic team located in Hershey, PA. This is an excellent opportunity to join an outstanding academic program with a national reputation and impact the lives of our future Emergency Medicine physicians.

### **What We're Offering:**

- Competitive salary and benefits
- Sign-On Bonus
- Relocation Assistance
- Leadership for Emergency Medicine Residency Program
- Comprehensive benefit and retirement options

### **What We're Seeking:**

- MD, DO, or foreign equivalent
- BC/BE by ABEM or ABOEM
- Leadership experience
- Outstanding patient care qualities
- Ability to work collaboratively within a diverse academic and clinical environment



### **FOR MORE INFORMATION PLEASE CONTACT:**

**Heather Peffley, PHR CPRP**  
Physician Recruiter  
Penn State Health

**Email:** [hpeffley@pennstatehealth.psu.edu](mailto:hpeffley@pennstatehealth.psu.edu)  
**Website:** [careers.pennstatehealth.org](https://careers.pennstatehealth.org)

### **What the Area Offers:**

Located in a safe family-friendly setting, Hershey, PA, our local neighborhoods boast a reasonable cost of living whether you prefer a more suburban setting or thriving city rich in theater, arts, and culture. Known as the home of the Hershey chocolate bar, Hershey's community is rich in history and offers an abundant range of outdoor activities, arts, and diverse experiences. We're conveniently located within a short distance to major cities such as Philadelphia, Pittsburgh, NYC, Baltimore, and Washington DC.

## THE BRASS TACKS: CONCISE REVIEWS OF PUBLISHED EVIDENCE

# Accuracy of the European Society of Cardiology 0/1-, 0/2-, and 0/3-hour algorithms for diagnosing acute myocardial infarction

Brit Long MD<sup>1</sup>  | Michael Gottlieb MD<sup>2</sup> 

<sup>1</sup>Department of Emergency Medicine, Brooke Army Medical Center, Fort Sam Houston, Texas, USA

<sup>2</sup>Department of Emergency Medicine, Rush University Medical Center, Chicago, Illinois, USA

### Correspondence

Brit Long, MD, Department of Emergency Medicine, Brooke Army Medical Center, 3841 Roger Brooke Drive, Fort Sam Houston, TX 78234, USA.  
Email: brit.long@yahoo.com

**Editor's Note:** Brass Tacks are concise reviews of published evidence. This series is a result of collaboration between *Academic Emergency Medicine* and the evidence-based medicine website [www.TheNNT.com](http://www.TheNNT.com). For inquiries please contact the section editor, Shahriar Zehtabchi, MD (email: [mshahriar.zehtabchi@downstate.edu](mailto:mshahriar.zehtabchi@downstate.edu)).

**Keywords:** cardiology, high-sensitivity troponin, myocardial infarction

Summary heading	The European Society of Cardiology algorithms have high sensitivity (0/1- and 0/2-h) and high specificity (0/1-, 0/2-, and 0/3-h) for AMI
Positive LR findings	LR+: 14 for 0/1-h algorithm for ruling in AMI LR+: 21 for 0/2-h LR+: 13 for 0/3-h
Negative LR findings	LR-: 0.01 for 0/1-h algorithm for ruling out AMI LR-: 0.02 for 0/2-h LR-: 0.09 for 0/3-h
Who was in the studies	32 studies comprising 30,066 patients with suspected AMI

## NARRATIVE

Chest pain is a common presentation to the emergency department (ED), representing over 5 million annual ED visits in the United States.<sup>1</sup> However, only 10%–20% of ED patients with chest pain are ultimately diagnosed with acute myocardial infarction (AMI).<sup>2</sup> Evaluation using accelerated diagnostic protocols with high-sensitivity cardiac troponin (hs-cTn) tests can provide more rapid detection of AMI and earlier discharge for patients in whom AMI has been excluded when compared to standard troponin assays.<sup>3,4</sup> Recent guidelines utilize hs-cTn in evaluation for AMI,<sup>5</sup> while prior guidelines incorporated standard troponin assays at the initial time of presentation and at 3 h.<sup>6–8</sup> The most recent European Society of Cardiology (ESC) guidelines use specific hs-cTn T or I thresholds at

0 h and 1 or 2 h and absolute changes to determine who may be ruled in or ruled out for AMI as well as those who require observation.<sup>5</sup>

The systematic review and meta-analysis discussed here included prospective observational cohort studies, implementation studies, and randomized controlled trials (RCTs) evaluating ESC hs-cTn protocols. Studies included adult patients in the ED or chest pain unit with suspected non-ST elevation myocardial infarction or acute coronary syndrome.<sup>9</sup> The authors only included studies evaluating the 0/1-, 0/2-, and 0/3-h ESC protocols utilizing the Elecsys hs-cTnT (Roche), Architect hs-cTnI (Abbott), and Centaur/Atellica hs-cTnI (Siemens) assays based on 2015 ESC guideline thresholds.<sup>7</sup> Diagnosis of AMI was determined based on the Third or Fourth Global Task Force Universal Definition of Myocardial Infarction.<sup>10,11</sup> The primary outcome was diagnostic accuracy for AMI using the 0/1-, 0/2-, and 0/3-h protocols.

The systematic review identified 32 publications ( $n = 30,066$  patients; 4246 cases of AMI) conducted in Asia, Australasia, Europe, and North America that met the inclusion criteria.<sup>9</sup> Among these 32 papers, they identified 33 total cohorts, of which 20 were unique subgroups. Sixteen cohorts evaluated the 0/1-h algorithm, seven cohorts the 0/2-h algorithm, and 10 cohorts the 0/3-h algorithm. Eighteen subgroups were evaluated using an observational study type, while one was evaluated with an RCT and one was quasi-experimental. The authors obtained primary data from the principal investigators for 16 subgroups and aggregate data from published articles for four subgroups. The prevalence of AMI ranged from 4% to 37% in the included studies.

The 0/1-h algorithm demonstrated high diagnostic accuracy to rule in (specificity = 94.0%, 95% confidence interval [CI] =

Supervising Editor: Shahriar Zehtabchi, MD.

90.7%–96.2%; positive likelihood ratio [LR+] = 14, 95% CI = 9–20) and to rule out AMI (sensitivity = 99%, 95% CI = 98.5%–99.5%; negative likelihood ratio [LR–] = 0.01, 95% CI = 0.01–0.02), with 17% of patients ruled in and 54% of patients ruled out. The 0/2-h algorithm demonstrated similar test characteristics, with high accuracy for ruling in (specificity = 96%, 95% CI = 92.9%–97.9%; LR+ = 21, 95% CI = 13–35) and ruling out AMI (sensitivity = 98.6%, 95% CI = 97.2%–99.3%; LR– = 0.02, 95% CI = 0.01–0.04), with 15% of patients ruled in and 61% of patients ruled out. The 0/3-h algorithm had similar ability to rule in AMI (specificity = 93%, 95% CI = 86.9%–96.6%; LR+ = 13, 95% CI = 6.7–24), but lower ability to rule out AMI (sensitivity = 93.7%, 95% CI = 87.4%–97.0%; LR– = 0.09, 95% CI = 0.05–0.15), with 19% of patients ruled in and 66% of patients ruled out. The proportion of patients who remained undifferentiated or in the observational zone was 29% for the 0/1-h algorithm, 24% for the 0/2-h algorithm, and 15% for the 0/3-h algorithm. Stratification by assay demonstrated similar sensitivities and specificities.

## CAVEATS

This study has several important limitations. While all three ESC algorithms had high specificities, there was significant heterogeneity across the included studies. This heterogeneity may be in part due to variation in AMI prevalence. Populations with a higher prevalence of AMI may also have a higher prevalence of non-AMI conditions associated with troponin elevations. The heterogeneity could also be due to differences in inclusion and exclusion criteria used by different studies, with studies including only patients with chest pain having higher specificities compared to studies which included patients presenting with other symptoms. There were significant differences in implementation of the 0/3-h protocols among the included studies as well as potential miscalibration of the Elecsys hs-cTnT lots used globally from 2010 to 2012, further contributing to the heterogeneity. The lower sensitivity of the 0/3-h protocol and wide CIs could be due to the fact that some studies used this algorithm without clinical criteria (GRACE score <140 and pain-free) while others studies used it combined with clinical criteria (sensitivity without and with clinical criteria = 90% [95% CI = 82.9%–94.6%] vs. 98% [95% CI = 88.6%–99.8%], respectively). In fact, protocol performance may be suboptimal in some populations such as those at high-risk for AMI clinically.<sup>12</sup> The performance of algorithms for diagnosis of AMI using sex-specific 99th percentile thresholds are still unclear as many studies did not adjudicate based on this factor. Several studies used samples collected in a different timing manner than that recommended in ESC guidelines, such as using a 0–2/3-h blood draw instead of a 0/2- or 0/3-h draw,<sup>13</sup> and the systematic review authors included studies analyzing troponin samples collected over 30 min outside of the stipulated time by the ESC algorithms. The authors were unable to obtain individual patient-level data and could not account for patients falsely ruled

in or out by the ESC algorithms. While most of these studies used the third universal definition of AMI, the adjudication criteria in each of these studies likely means that the difference between third and fourth universal definitions are unlikely to substantially bias the results. Finally, patient management was not dictated by the algorithms and, therefore, outcomes may have been influenced by troponin levels and clinical management.

## CONFLICT OF INTEREST

The authors declare no potential conflict of interest.

## ORCID

Brit Long  <https://orcid.org/0000-0003-4770-8869>

Michael Gottlieb  <https://orcid.org/0000-0003-3276-8375>

## REFERENCES

1. Aalam AA, Alsabban A, Pines JM. National trends in chest pain visits in US emergency departments (2006–2016). *Emerg Med J*. 2020;37(11):696–699.
2. Goodacre S, Cross E, Arnold J, Angelini K, Capewell S, Nicholl J. The health care burden of acute chest pain. *Heart*. 2005;91(2):229–230.
3. Eggers KM, Lindahl B, Melki D, Jernberg T. Consequences of implementing a cardiac troponin assay with improved sensitivity at Swedish coronary care units: an analysis from the SWEDHEART registry. *Eur Heart J*. 2016;37(30):2417–2424.
4. Nejatian A, Omstedt Å, Höijer J, et al. Outcomes in patients with chest pain discharged after evaluation using a high-sensitivity troponin T assay. *J Am Coll Cardiol*. 2017;69(21):2622–2630.
5. Collet JP, Thiele H, Barbato E, et al. 2020 ESC guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Eur Heart J*. 2021;42(14):1289–1367. doi:10.1093/eurheartj/ehaa575. Erratum in: *Eur Heart J*. 2021 May 14;42(19):1908. Erratum in: *Eur Heart J*. 2021 May 14;42(19):1925. Erratum in: *Eur Heart J*. 2021 May 13; PMID: 32860058.
6. Hamm CW, Bassand JP, Agewall S, et al. ESC guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: the Task Force for the Management of Acute Coronary Syndromes (ACS) in Patients Presenting Without Persistent ST-segment Elevation of the European Society of Cardiology (ESC). *Eur Heart J*. 2011;32(23):2999–3054.
7. Anand A, Shah AS, Beshiri A, Jaffe AS, Mills NL. Global adoption of high-sensitivity cardiac troponins and the universal definition of myocardial infarction. *Clin Chem*. 2019;65(3):484–489.
8. Roffi M, Patrono C, Collet JP, et al. 2015 ESC guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: Task Force for the Management of Acute Coronary Syndromes in Patients Presenting Without Persistent ST-Segment Elevation of the European Society of Cardiology (ESC). *Eur Heart J*. 2016;37(3):267–315.
9. Chiang CH, Chiang CH, Pickering JW, et al. Performance of the European Society of Cardiology 0/1-hour, 0/2-hour, and 0/3-hour algorithms for rapid triage of acute myocardial infarction: an international collaborative meta-analysis. *Ann Intern Med*. 2022;175(1):101–113.
10. Thygesen K, Alpert JS, Jaffe AS, et al. Third universal definition of myocardial infarction. *Circulation*. 2012;126(16):2020–2035.
11. Thygesen K, Alpert JS, Jaffe AS, et al. Fourth universal definition of myocardial infarction (2018). *Circulation*. 2018;138(20):e618–e651. doi:10.1161/CIR.0000000000000617. Erratum in: *Circulation*. 2018 Nov 13;138(20):e652.

12. Kavsak PA, Hewitt MK, Mondoux SE, et al. Diagnostic performance of serial high-sensitivity cardiac troponin measurements in the emergency setting. *J Cardiovasc Dev Dis.* 2021;8(8):97.
13. Nowak RM, Christenson RH, Jacobsen G, et al. Performance of novel high-sensitivity cardiac troponin I assays for 0/1-hour and 0/2- to 3-hour evaluations for acute myocardial infarction: results from the HIGH-US study. *Ann Emerg Med.* 2020;76(1):1-13.

**How to cite this article:** Long B & Gottlieb M. Accuracy of the European Society of Cardiology 0/1-, 0/2-, and 0/3-hour algorithms for diagnosing acute myocardial infarction. *Acad Emerg Med.* 2022;00:1-3. doi: [10.1111/acem.14444](https://doi.org/10.1111/acem.14444)