

Lumbar Puncture After Negative Imaging in Patients With Suspected Subarachnoid Hemorrhage

Opposing authors provide succinct, authoritative discussions of controversial issues in emergency medicine. Authors are provided the opportunity to review and comment on opposing presentations. Each topic is accompanied by an Editor's Note that summarizes important concepts. Participation as an authoritative discussant is by invitation only, but suggestions for topics and potential authors can be submitted to the section editors.

Editor's Note: The sensitivity of computed tomographic (CT) head imaging in detecting subarachnoid hemorrhage has substantially increased in recent years. As a consequence, very few patients with a negative CT scan result will actually have subarachnoid hemorrhage. Efforts to identify patients with rare subarachnoid hemorrhage will require performing lumbar punctures on a large population of disease-free individuals. In this installment of Clinical Controversies, opposing advocates address the issues that arise in subjecting a large population to unpleasant evaluations to detect a dangerous diagnosis in a small minority.

LUMBAR PUNCTURE IS NECESSARY FOR RULING OUT ATRAUMATIC SUBARACHNOID HEMORRHAGE SIX HOURS AFTER SYMPTOM ONSET



Michael D. April, MD, DPhil
2nd Brigade Combat Team, 4th Infantry Division
Fort Carson, CO; and
Department of Military and Emergency Medicine
Uniformed Services University of the Health Sciences
Bethesda, MD

Brit Long, MD
Department of Emergency Medicine
San Antonio Uniformed Services Health Education Consortium
Fort Sam Houston, TX

[Ann Emerg Med. 2021;77:641-643.]

Subarachnoid hemorrhage is a deadly disease that commonly presents with acute onset of atraumatic headache. Initial misdiagnosis leads to worse outcomes owing to delays in management.¹ High-risk patients with negative computed tomographic (CT) imaging results

obtained more than 6 hours from symptom onset should undergo lumbar puncture as opposed to no further evaluation.

Noncontrast CT of the head followed by lumbar puncture if the result is negative is the historical mainstay for the diagnosis of atraumatic subarachnoid hemorrhage. Yet CT sensitivity decreases with time from symptom onset. A prospective cohort study of 3,132 patients reported a sensitivity of 100% within 6 hours of symptom onset but 85.7% thereafter.² Meta-analysis data suggest that a negative CT result obtained greater than 6 hours from symptom onset yields a negative likelihood ratio (LR-) of 0.07 for the diagnosis of subarachnoid hemorrhage.³

Adding lumbar puncture to CT imaging yields a near 100% sensitivity for the diagnosis of subarachnoid hemorrhage regardless of time elapsed from symptom onset.⁴ Estimated lumbar puncture numbers needed to identify actionable cases of subarachnoid hemorrhage are highly situation dependent. Published estimates range from 91 to 15,200.³ We believe these numbers represent overestimates because they come from studies examining lumbar punctures for any indication or studies excluding patients with features placing them at high risk for subarachnoid hemorrhage. Lumbar puncture offers the most sensitive tool for ruling out subarachnoid hemorrhage and also facilitates the diagnosis of alternative actionable diagnoses such as meningitis.⁴

Lumbar punctures are not without risk. Potential complications include post-lumbar puncture headache, bleeding, and brainstem herniation. That said, the risk of catastrophic complications such as herniation is low, provided clinicians appropriately screen patients for

contraindications such as bleeding diatheses and focal neurologic deficits.³ Unfortunately, traumatic taps are common, which can erroneously inflate the numbers of RBCs in the cerebrospinal fluid. Because this type of tap occurs in up to 15% of procedures, results can be challenging to interpret when clinicians are trying to avoid false-positive ones.⁵ The resultant loss in specificity is the price clinicians pay for the sensitivity offered by this procedure.

Whether a particular patient is more likely to experience benefit versus harm from a lumbar puncture procedure depends entirely on the probability of subarachnoid hemorrhage in that patient. Historical and physical examination features define probability of disease, in turn driving the utility of diagnostic procedures.³ The Ottawa Subarachnoid Hemorrhage Rule defines characteristics of patients at particularly high risk for subarachnoid hemorrhage: older than 40 years, complaint of neck pain or stiffness, witnessed loss of consciousness, onset with exertion, thunderclap headache, and limited neck flexion on examination.^{6,7} Patients at highest risk of subarachnoid hemorrhage are most likely to benefit from lumbar puncture.

To illustrate this point, consider 3 separate patients. First is an intermediate-risk patient whose risk of subarachnoid hemorrhage matches the prevalence identified in a meta-analysis of studies of subarachnoid hemorrhage diagnostic evaluation, 7.5%.³ Second is a low-risk patient whose headache did not reach maximal intensity within 1 hour, a feature with an LR⁻ of 0.06, yielding a subarachnoid hemorrhage probability of 4.8%. Third is a high-risk patient with neck stiffness on examination, a feature with a positive LR of 6.6, yielding a subarachnoid hemorrhage probability of 34.8%. Given an LR⁻ of 0.07, a negative CT result obtained greater than 6 hours from symptom onset yields a posterior probability of disease of 0.3% for the low-risk patient, 0.6% for the intermediate-risk patient, and 2.3% for the high-risk patient (Figure). Using Pauker's formula for testing threshold,⁸ we assume lumbar puncture sensitivity is 100%,⁴ specificity is 67%,⁴ and risk of mortality approximates 0%. We assume mortality reduction owing to surgical intervention is 14%¹ and periprocedural mortality is 0.7%.⁹ This yields a testing threshold of 1.6%. The high-risk patient's posterior probability of




Figure. Fagan nomogram. The left axis represents the pretest probability of subarachnoid hemorrhage. The center axis represents the LR of a negative result for noncontrast CT of the brain obtained greater than 6 hours from symptom onset (0.07). The right axis represents the posttest probability of subarachnoid hemorrhage after the negative CT result. In the case of a low-risk patient with a pretest probability of subarachnoid hemorrhage of 4.8%, the posttest probability of subarachnoid hemorrhage is 0.3% (dashed line). For an intermediate-risk patient with a pretest probability of subarachnoid hemorrhage of 7.5%, posttest subarachnoid hemorrhage probability decreases to 0.6% (solid line). A high-risk patient with a pretest probability of disease of 34.8% still retains a posttest probability of subarachnoid hemorrhage of 2.3% (bold line).

disease exceeds this value; hence, he or she should undergo lumbar puncture. For the other 2 patients, lumbar puncture is likely to do more harm than good.

Clinical reality is that scenarios will arise in which the probability of disease approximates the testing threshold, leading to decisional equipoise. Shared decisionmaking can balance the risks and benefits in determining diagnostic strategy in these patients. Subarachnoid hemorrhage represents a dangerous condition, the presentation of which is often not distinct from that of the high volume of emergency department patients with headache. Hence, the diagnostic evaluation for this condition is not always clear, making it an optimal scenario for use of shared decisionmaking.¹⁰ Yet among high-risk patients with negative-result head CT imaging obtained greater than 6 hours from symptom onset, we believe lumbar puncture must be the default course of action to minimize the risk of a missed subarachnoid hemorrhage diagnosis.

This review does not reflect the views or opinions of the US government, Department of Defense, US Army, US Air Force, Brooke Army Medical Center, or SAUSHEC EM Residency Program.

<https://doi.org/10.1016/j.annemergmed.2020.11.009>

REFERENCES

1. Kowalski RG, Claassen J, Kreiter KT, et al. Initial misdiagnosis and outcome after subarachnoid hemorrhage. *JAMA*. 2004;291:866-869.
2. Perry JJ, Stiell IG, Sivilotti ML, et al. Sensitivity of computed tomography performed within six hours of onset of headache for diagnosis of subarachnoid haemorrhage: prospective cohort study. *BMJ*. 2011;343:d4277.
3. Carpenter CR, Hussain AM, Ward MJ, et al. Spontaneous subarachnoid hemorrhage: a systematic review and meta-analysis describing the diagnostic accuracy of history, physical examination, imaging, and lumbar puncture with an exploration of test thresholds. *Acad Emerg Med*. 2016;23:963-1003.
4. Perry JJ, Spacek A, Forbes M, et al. Is the combination of negative computed tomography result and negative lumbar puncture result sufficient to rule out subarachnoid hemorrhage? *Ann Emerg Med*. 2008;51:707-713.
5. Shah KH, Richard KM, Nicholas S, et al. Incidence of traumatic lumbar puncture. *Acad Emerg Med*. 2003;10:151-154.
6. Perry JJ, Sivilotti MLA, Emond M, et al. Prospective implementation of the Ottawa Subarachnoid Hemorrhage Rule and 6-Hour Computed Tomography Rule. *Stroke*. 2020;51:424-430.
7. American College of Emergency Physicians Clinical Policies Subcommittee (Writing Committee) on Acute Headache. Clinical policy: critical issues in the evaluation and management of adult patients presenting to the emergency department with acute headache. *Ann Emerg Med*. 2019;74:e41-e74.
8. Pauker SG, Kassirer JP. The threshold approach to clinical decision making. *N Engl J Med*. 1980;302:1109-1117.
9. Hwang JS, Hyun MK, Lee HJ, et al. Endovascular coiling versus neurosurgical clipping in patients with unruptured intracranial aneurysm: a systematic review. *BMC Neurol*. 2012;12:99.
10. Hess EP, Grudzen CR, Thomson R, et al. Shared decision-making in the emergency department: respecting patient autonomy when seconds count. *Acad Emerg Med*. 2015;22:856-864.

LUMBAR PUNCTURE SHOULD NOT BE ROUTINELY PERFORMED FOR SUBARACHNOID HEMORRHAGE AFTER A NEGATIVE HEAD CT

Michael Gottlieb, MD

Department of Emergency Medicine
Rush University Medical Center
Chicago, IL

Justin Morgenstern, MD

Division of Emergency Medicine
Department of Family and Community Medicine
University of Toronto
Toronto, Ontario, Canada

[Ann Emerg Med. 2021;77:643-645.]

Headache is a common presenting complaint in the emergency department (ED), and approximately 1% of patients with a sudden-onset headache receive a diagnosis of subarachnoid hemorrhage.¹ Historically, it has been taught that a lumbar puncture is required after a negative computed tomographic (CT) result to exclude the diagnosis. However, in light of improving CT quality, the value of this approach has been questioned. In accordance with the current evidence, we argue that lumbar puncture is not routinely required to rule out subarachnoid hemorrhage after a negative head CT scan result, even if the CT was performed more than 6 hours after symptom onset.

The most important argument against lumbar puncture is the accuracy of modern CT scanners. The tradition of performing a lumbar puncture after CT started in an era when CT was more likely to miss a diagnosis of subarachnoid hemorrhage. Using modern CT technology, Perry et al² demonstrated 100% sensitivity (95% confidence interval 97% to 100%) of CT within 6 hours for the diagnosis of subarachnoid hemorrhage.² Although a CT performed greater than 6 hours after symptom onset is less sensitive, the prevalence of subarachnoid hemorrhage also decreases among patients presenting later.³ In the original study by Perry et al,² there were 2,179 patients

